

MULTIFRACTAL ANALYSIS OF THE DIVERGENCE OF FOURIER SERIES

FRÉDÉRIC BAYART, YANICK HEURTEAUX

ABSTRACT. A famous theorem of Carleson says that, given any function $f \in L^p(\mathbb{T})$, $p \in (1, +\infty)$, its Fourier series $(S_n f(x))$ converges for almost every $x \in \mathbb{T}$. Beside this property, the series may diverge at some point, without exceeding $O(n^{1/p})$. We define the divergence index at x as the infimum of the positive real numbers β such that $S_n f(x) = O(n^\beta)$ and we are interested in the size of the exceptional sets E_β , namely the sets of $x \in \mathbb{T}$ with divergence index equal to β . We show that quasi-all functions in $L^p(\mathbb{T})$ have a multifractal behavior with respect to this definition. Precisely, for quasi-all functions in $L^p(\mathbb{T})$, for all $\beta \in [0, 1/p]$, E_β has Hausdorff dimension equal to $1 - \beta p$. We also investigate the same problem in $C(\mathbb{T})$, replacing polynomial divergence by logarithmic divergence. In this context, the results that we get on the size of the exceptional sets are rather surprising.

1. INTRODUCTION

1.1. Description of the results. The famous theorem of Carleson and Hunt asserts that, when f belongs to $L^p(\mathbb{T})$, $1 < p < +\infty$, where $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, the sequence of the partial sums of its Fourier series $(S_n f(x))_{n \geq 0}$ converges for almost every $x \in \mathbb{T}$. On the other hand, it can diverge at some point. This divergence cannot be too fast since, for any $f \in L^p(\mathbb{T})$ and any $x \in \mathbb{T}$, $|S_n f(x)| \leq C_p n^{1/p} \|f\|_p$ (see [11] for instance). In view of these results, a natural question arises. How big can be the sets F such that $|S_n f(x)|$ grows as fast as possible for every $x \in F$? More generally, can we say something on the size of the sets such that $|S_n f(x)|$ behaves like (or as bad as) n^β for some $\beta \in (0, 1/p]$?

To measure the size of subsets of \mathbb{T} , we shall use the Hausdorff dimension. Let us recall the relevant definitions (we refer to [5] and to [10] for more on this subject). If $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a nondecreasing continuous function satisfying $\phi(0) = 0$ (ϕ is called a *dimension function* or a *jauge function*), the ϕ -Hausdorff outer measure of a set $E \subset \mathbb{R}^d$ is

$$\mathcal{H}^\phi(E) = \lim_{\varepsilon \rightarrow 0} \inf_{r \in R_\varepsilon(E)} \sum_{B \in r} \phi(|B|),$$

$R_\varepsilon(E)$ being the set of countable coverings of E with balls B of diameter $|B| \leq \varepsilon$. When $\phi_s(x) = x^s$, we write for short \mathcal{H}^s instead of \mathcal{H}^{ϕ_s} . The Hausdorff dimension of a set E is

$$\dim_{\mathcal{H}}(E) := \sup\{s > 0; \mathcal{H}^s(E) > 0\} = \inf\{s > 0; \mathcal{H}^s(E) = 0\}.$$

The first result studying the Hausdorff dimension of the divergence sets of Fourier series is due to J-M. Aubry [2].

Date: March 2, 2011.

Theorem 1.1. *Let $f \in L^p(\mathbb{T})$, $1 < p < +\infty$. For $\beta \geq 0$, define*

$$\mathcal{E}(\beta, f) = \left\{ x \in \mathbb{T}; \limsup_{n \rightarrow +\infty} n^{-\beta} |S_n f(x)| > 0 \right\}.$$

Then $\dim_{\mathcal{H}}(\mathcal{E}(\beta, f)) \leq 1 - \beta p$. Conversely, given a set E such that $\dim_{\mathcal{H}}(E) < 1 - \beta p$, there exists a function $f \in L^p(\mathbb{T})$ such that, for any $x \in E$, $\limsup_{n \rightarrow +\infty} n^{-\beta} |S_n f(x)| = +\infty$.

This result motivated us to introduce the notion of divergence index. For a given function $f \in L^p(\mathbb{T})$ and a given point $x_0 \in \mathbb{T}$, we can define the real number $\beta(x_0)$ as the infimum of the non negative real numbers β such that $|S_n f(x_0)| = O(n^\beta)$. The real number $\beta(x_0)$ will be called the *divergence index* of the Fourier series of f at point x_0 . Of course, for any function $f \in L^p(\mathbb{T})$ ($1 < p < +\infty$) and any point $x_0 \in \mathbb{T}$, $0 \leq \beta(x_0) \leq 1/p$. Moreover, Carleson's theorem implies that $\beta(x_0) = 0$ almost surely and we would like to have precise estimates on the size of the level sets of the function β . These are defined as

$$\begin{aligned} E(\beta, f) &= \{x \in \mathbb{T}; \beta(x) = \beta\} \\ &= \left\{ x \in \mathbb{T}; \limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log n} = \beta \right\}. \end{aligned}$$

We can ask for which values of β the sets $E(\beta, f)$ are non-empty. This set of values will be called the domain of definition of the spectrum of singularities of f . If β belongs to the domain of definition of the spectrum of singularities, it is also interesting to estimate the Hausdorff dimension of the sets $E(\beta, f)$. The function $\beta \mapsto \dim_{\mathcal{H}}(E(\beta, f))$ will be called the spectrum of singularities of the function f (in terms of its Fourier series). By Aubry's result, $\dim_{\mathcal{H}}(E(\beta, f)) \leq 1 - \beta p$ and, for any fixed $\beta_0 \in [0, 1/p]$, for any $\varepsilon > 0$, one can find $f \in L^p(\mathbb{T})$ such that $\dim_{\mathcal{H}}\left(\bigcup_{\beta_0 \leq \beta \leq 1/p} E(\beta, f)\right) \geq 1 - \beta_0 p - \varepsilon$. Our first main result is that a *typical* function $f \in L^p(\mathbb{T})$ satisfies $\dim_{\mathcal{H}}(E(\beta, f)) = 1 - \beta p$ for *any* $\beta \in [0, 1/p]$. In particular, f has a multifractal behavior with respect to the summation of its Fourier series, meaning that the domain of definition of its spectrum of singularities contains an interval with non-empty interior.

Theorem 1.2. *Let $1 < p < +\infty$. For quasi-all functions $f \in L^p(\mathbb{T})$, for any $\beta \in [0, 1/p]$, $\dim_{\mathcal{H}}(E(\beta, f)) = 1 - \beta p$.*

The terminology "quasi-all" used here is relative to the Baire category theorem. It means that this property is true for a residual set of functions in $L^p(\mathbb{T})$.

In a second part of the paper, we turn to the case of $\mathcal{C}(\mathbb{T})$, the set of continuous functions on \mathbb{T} . In that space, the divergence of Fourier series is controlled by a logarithmic factor. More precisely, if (D_n) is the sequence of the Dirichlet kernels, we know that $\|S_n f\|_\infty \leq \|D_n\|_1 \|f\|_\infty$, so that there exists some absolute constant $C > 0$ such that $\|S_n f\|_\infty \leq C \|f\|_\infty \log n$ for any $f \in \mathcal{C}(\mathbb{T})$ and any $n > 1$. As before, one can discuss the size of the sets such that $|S_n f(x)|$ behaves badly, namely like $(\log n)^\beta$, $\beta \in [0, 1]$. More precisely, mimicking the case of the L^p spaces, we introduce, for any $\beta \in [0, 1]$ and any $f \in \mathcal{C}(\mathbb{T})$,

the following sets:

$$\begin{aligned}\mathcal{F}(\beta, f) &= \left\{ x \in \mathbb{T}; \limsup_{n \rightarrow +\infty} (\log n)^{-\beta} |S_n f(x)| > 0 \right\} \\ F(\beta, f) &= \left\{ x \in \mathbb{T}; \limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log \log n} = \beta \right\}.\end{aligned}$$

Theorem 1.1 indicates that, on $L^p(\mathbb{T})$, $|S_n f(x)|$ can grow as fast as possible (namely like $n^{1/p}$) only on small sets: for every function $f \in L^p(\mathbb{T})$, $\dim_{\mathcal{H}}(E(1/p, f)) = 0$. This property dramatically breaks down on $\mathcal{C}(\mathbb{T})$, as the following result indicates.

Theorem 1.3. *For quasi-all functions $f \in \mathcal{C}(\mathbb{T})$, $\dim_{\mathcal{H}}(F(1, f)) = 1$.*

Thus, for quasi-all functions $f \in \mathcal{C}(\mathbb{T})$, the partial sums $(S_n f(x))_{n \geq 0}$ grow as fast as possible on big sets.

We can also study the domain of the spectrum of singularities of f , namely the values of β such that $F(\beta, f)$ is non-empty. Like in the case of the space $L^p(\mathbb{T})$, this domain is for quasi-all functions of $\mathcal{C}(\mathbb{T})$ an interval with non-empty interior, so that a typical function f in $\mathcal{C}(\mathbb{T})$ has a multifractal behavior with respect to the summation of its Fourier series. However, the spectrum of singularities is constant!

Theorem 1.4. *For quasi-all functions $f \in \mathcal{C}(\mathbb{T})$, for any $\beta \in [0, 1]$, $F(\beta, f)$ is non-empty and has Hausdorff dimension 1.*

Theorem 1.4 indicates that the Hausdorff dimension is not precise enough to measure the size of the level sets $F(\beta, f)$. This leads us to introduce a notion of *precised Hausdorff dimension*, in order to distinguish more finely sets which have the same Hausdorff dimension. For $s > 0$ and $t \in (0, 1]$, we consider

$$\phi_{s,t}(x) = x^s \exp [(\log 1/x)^{1-t}].$$

Definition 1.5. Let $E \subset \mathbb{R}^d$. We say that E has *precised Hausdorff dimension* (α, β) if α is the Hausdorff dimension of E and

- $\beta = 0$ if $\mathcal{H}^{\phi_{\alpha,t}}(E) = 0$ for every $t \in (0, 1)$;
- $\beta = \sup \{t \in (0, 1); \mathcal{H}^{\phi_{\alpha,t}}(E) > 0\}$ otherwise.

It is not difficult to check that $\phi_{s,t}(x) \leq \phi_{s',t'}(x)$ for small values of x iff

$$s > s' \text{ or } (s = s' \text{ and } t \geq t').$$

Thus the precised Hausdorff dimension is a refinement of the Hausdorff dimension. In particular it is a tool to classify sets that have the same Hausdorff dimension. The natural order for the precised dimension (s, t) is the lexicographical order which will be denoted by \prec . With respect to this order, we can say that the greater is the set, the greater is the precised dimension. Moreover, if $(s, t) \prec (s', t')$ and $(s, t) \neq (s', t')$, then $\phi_{s',t'} \ll \phi_{s,t}$. It follows that $H^{\phi_{s',t'}}(E) = 0$ as soon as $H^{\phi_{s,t}}(E) < \infty$.

Our main theorem on $\mathcal{C}(\mathbb{T})$, which contains both Theorems 1.3 and 1.4, is the following:

Theorem 1.6. *For quasi-all functions $f \in \mathcal{C}(\mathbb{T})$, for any $\beta \in [0, 1]$, the precised Hausdorff dimension of $F(\beta, f)$ is $(1, 1 - \beta)$.*

The paper is organized as follows. In the remaining part of this section, we introduce tools which will be needed during the rest of the paper. In Section 2, we prove Theorem 1.2 whereas in Section 3, we prove Theorem 1.6.

1.2. A precised version of Fejér's theorem. Working on Fourier series, we will need results on approximation by trigonometric polynomials. Let $k \in \mathbb{Z}$ and $e_k : t \mapsto e^{2\pi ikt}$, so that, for any $g \in L^1(\mathbb{T})$ and any $n \in \mathbb{N}$,

$$S_n g : t \mapsto \sum_{k=-n}^n \langle g, e_k \rangle e_k(t).$$

Let $\sigma_n g$ be the n -th Fejér sum of g ,

$$\sigma_n g : t \mapsto \frac{1}{n} \sum_{k=0}^{n-1} S_k g(t).$$

$\sigma_n g$ is obtained by taking the convolution of g with the Fejér kernel

$$F_n : t \mapsto \frac{1}{n} \left(\frac{\sin(n\pi t)}{\sin(\pi t)} \right)^2.$$

If g belongs to $\mathcal{C}(\mathbb{T})$, $(\sigma_n g)_{n \geq 1}$ converges uniformly to g . For our purpose, we need to estimate how quick is the convergence. This is the content of the next lemma (part (1) rectifies a mistake in the proof of Lemma 12 in [2] and requires to replace $\|\theta\|_\infty/4$ in Aubry's version by $\|\theta\|_\infty/2$).

Lemma 1.7. *Let θ be a Lipschitz function on \mathbb{T} , let $n \in \mathbb{N}$ and let $x \in \mathbb{T}$. Suppose that $\|\theta'\|_\infty \leq n$ and that $\theta(x) = 0$. Then the two following inequalities hold:*

$$(1) \quad |\sigma_n \theta(x)| \leq \frac{1}{4} + \frac{1}{2} \|\theta\|_\infty \quad \text{for any } n \geq 8$$

$$(2) \quad |\sigma_n \theta(x)| \leq 4 + \frac{1}{4} \|\theta\|_\infty \quad \text{for any } n \geq 4.$$

Proof. We may assume that $x = 0$. Hence, $\sigma_n \theta(0) = \int_{-1/2}^{1/2} \theta(y) F_n(y) dy$. Let us consider $\delta \in (0, 2]$ and $n \geq 4$. On the one hand, for any $y \in [0, 1/2)$,

$$0 \leq F_n(y) = \frac{\sin^2(n\pi y)}{n \sin^2(\pi y)} \leq \frac{1}{n(2y)^2}$$

so that

$$\left| \int_{\delta/n < |y| \leq 1/2} \theta(y) F_n(y) dy \right| \leq \frac{1}{2n} \|\theta\|_\infty \int_{\delta/n}^{+\infty} \frac{dy}{y^2} = \frac{\|\theta\|_\infty}{2\delta}.$$

On the other hand,

$$\left| \int_{-\delta/n}^{\delta/n} \theta(y) F_n(y) dy \right| \leq 2 \int_0^{\delta/n} \left(\frac{\sin(n\pi y)}{\sin(\pi y)} \right)^2 y dy := u_n.$$

Using the convexity inequality $\sin\left(\frac{n}{n+1}\pi y\right) \geq \frac{n}{n+1} \sin(\pi y)$ and a change of variables, we see that (u_n) is non-increasing. To prove (1), we choose $\delta = 1$ and we observe that

$u_8 = 0.2496\dots \leq \frac{1}{4}$. To prove (2), we choose $\delta = 2$ and we observe that, since the maximum of F_n is $F_n(0) = n$,

$$|u_n| \leq 2n^2 \int_0^{2/n} y dy = 4.$$

□

1.3. The mass transference principle. The second main tool that we need in this paper is a method to produce sets with large Hausdorff dimension (Theorem 1.2) or with large precised Hausdorff dimension (Theorem 1.6). An efficient way to do this is to consider *ubiquitous systems* like this was done in [4, 7]. This was later refined in [3] to obtain a general mass transference principle, which we recall in the form that we need.

Theorem 1.8 (The mass transference principle). *Let $(x_n)_{n \geq 0}$ be a sequence of points in $[0, 1]^d$ and let $(r_n)_{n \geq 0}$ be a sequence of positive real numbers decreasing to 0. Let also $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a dimension function satisfying $\phi(s) \gg s^d$ when s goes to 0 and $s^{-d}\phi$ is monotonic. Define*

$$\begin{aligned} E &= \limsup_n B(x_n, r_n) \\ E^\phi &= \limsup_n B(x_n, \phi^{-1}(r_n^d)) \end{aligned}$$

and suppose that almost every point of $[0, 1]^d$ (in the sense of the Lebesgue's measure) lies in E . Then, $\mathcal{H}^\phi(E^\phi) = +\infty$.

We shall use it in the following situation.

Corollary 1.9. *Let (q_n) be a sequence of integers and, for each $n \in \mathbb{N}$, each $k \leq q_n$, let $B_{k,n} = B(x_{k,n}, r_{k,n})$ be a ball with center $x_{k,n} \in [0, 1]^d$ and with radius $r_{k,n}$ such that $\lim_{n \rightarrow +\infty} \max_k(r_{k,n}) = 0$. Let also $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a dimension function satisfying $\phi(s) \gg s^d$ when s goes to 0 and $s^{-d}\phi$ is monotonic. Define*

$$\begin{aligned} B_n &= \bigcup_{k=1}^{q_n} B_{k,n} & E &= \limsup_n B_n \\ B_n^\phi &= \bigcup_{k=1}^{q_n} B(x_{k,n}, \phi^{-1}(r_{k,n}^d)) & E^\phi &= \limsup_n B_n^\phi. \end{aligned}$$

Suppose that almost every point of $[0, 1]^d$ (in the sense of the Lebesgue's measure) lies in E . Then, $\mathcal{H}^\phi(E^\phi) = +\infty$.

Proof. Reordering the sequences $(B_{k,n})$ and $(B_{k,n}^\phi)$ as (C_j) and (C_j^ϕ) , we can observe that

$$\begin{aligned} \limsup_n B_n &= \limsup_j C_j = E \\ \limsup_n B_n^\phi &= \limsup_j C_j^\phi = E^\phi. \end{aligned}$$

Thus the corollary follows from a direct application of Theorem 1.8. □

2. MULTIFRACTAL ANALYSIS OF THE DIVERGENCE OF THE FOURIER SERIES OF
FUNCTIONS OF $L^p(\mathbb{T})$

In this section, we shall prove Theorem 1.2. Our method, which is inspired by [6], is divided into two parts. During the first one, we will construct a single function, which we call the saturating function, satisfying the conclusions of Theorem 1.2. During the second one, we will show how to derive a residual set from this single function.

2.1. The saturating function. Our intention is to construct a function g such that $|S_n g(x)|$ is big when x is close to a dyadic number. The following definition gives a precise meaning.

Definition 2.1. A real number x is α -approximable by dyadics, $\alpha \geq 1$, if there exist two sequence of integers $(k_n), (j_n)$ such that

$$\left| x - \frac{k_n}{2^{j_n}} \right| \leq \frac{1}{2^{\alpha j_n}}$$

and (j_n) goes to infinity. The dyadic exponent of x is the supremum of the set of real numbers α such that x is α -approximable by dyadics.

We denote by

$$D_\alpha = \{x \in [0, 1]; x \text{ is } \alpha\text{-approximable by dyadics}\}.$$

It is easy to check that $\mathcal{H}^\beta(D_\alpha) = 0$ for $\beta > 1/\alpha$ so that $\dim_{\mathcal{H}}(D_\alpha) \leq 1/\alpha$. On the other hand, it is well-known that $\dim_{\mathcal{H}}(D_\alpha) \geq \frac{1}{\alpha}$. Let us nevertheless show how this follows from Corollary 1.9. Indeed, D_α can be described as a limsup set:

$$D_\alpha = \limsup_{j \rightarrow +\infty} \bigcup_{k=0}^{2^j-1} I_{k,j}^\alpha$$

where the $I_{k,j}$ are the dyadic intervals

$$I_{k,j} = \left[\frac{k}{2^j} - \frac{1}{2^j}, \frac{k}{2^j} + \frac{1}{2^j} \right]$$

and

$$I_{k,j}^\alpha = \left[\frac{k}{2^j} - \frac{1}{2^{\alpha j}}, \frac{k}{2^j} + \frac{1}{2^{\alpha j}} \right].$$

Since $\bigcup_{k=0}^{2^j-1} I_{k,j} \supset [0, 1]$, Corollary 1.9 implies that $\mathcal{H}^{1/\alpha}(D_\alpha) = +\infty$.

We are going to define $g \in L^p(\mathbb{T})$ such that the divergence index of g at x depends on the dyadic exponent of x . The greater the dyadic exponent will be, the greater the divergence index of g at x will be. To do this, we will classify the dyadic intervals following their center. Namely, each $k/2^j$ can be uniquely written $K/2^J$ with $K \notin 2\mathbb{Z}$ and $1 \leq J \leq j$ (such a center comes into play from the J -th generation). Let $\mathcal{I}_J = \{K/2^J; K \notin 2\mathbb{Z}, 0 \leq K \leq 2^J - 1\}$ and

$$\mathbf{I}_{J,j} = \bigcup_{\frac{k}{2^j} \in \mathcal{I}_J} I_{k,j} \quad \mathbf{I}'_{J,j} = \bigcup_{\frac{k}{2^j} \in \mathcal{I}_J} 2I_{k,j}.$$

Here and elsewhere, when I is an interval and c is a positive real number, cI means the interval with the same center as I and with length $c|I|$. Observe that, when $1 \leq J < j$, the

intervals $2I_{k,j}$, $\frac{k}{2^j} \in \mathcal{I}_J$ don't overlap and the set $\mathbf{I}'_{J,j}$ has measure $2^{J-1}2^{2-j}$. Observe also that when J is small with respect to j , the real numbers x in $\mathbf{I}_{J,j}$ are well-approximated by dyadics $K/2^J$, since $|x - K/2^J| \leq 1/2^j$.

We first define a trigonometric polynomial with L^p -norm 1 which is almost constant on each $\mathcal{I}_{J,j}$ and which is big on $\mathcal{I}_{J,j}$ when J is small.

Lemma 2.2. *Let $j \geq 1$. There exists a trigonometric polynomial $g_j \in L^p(\mathbb{T})$ with spectrum contained in $[0, j2^{j+1})$ such that*

- $\|g_j\|_p \leq 1$;
- For any $1 \leq J \leq j$ and any $x \in \mathbf{I}_{J,j}$, we can find two integers n_1 and n_2 satisfying $0 \leq n_1 < n_2 < j2^{j+1}$ and such that

$$|S_{n_2}g_j(x) - S_{n_1}g_j(x)| \geq \frac{1}{4j}2^{-(J-j+1)/p}.$$

Proof. We set for any $1 \leq J \leq j$:

- $\chi_{J,j}$ a continuous piecewise linear function equal to 1 on $\mathbf{I}_{J,j}$, equal to 0 outside $\mathbf{I}'_{J,j}$, and satisfying $0 \leq \chi_{J,j} \leq 1$ and $\|\chi'_{J,j}\|_\infty \leq 2^j$;
- $c_{J,j} = \frac{1}{j}2^{-(J-j+1)/p}$ ($c_{J,j}$ is big when J is small);
- $g_{J,j} = e^{(2J-1)2^j\sigma_{2^j}}\chi_{J,j}$.

It is straightforward to observe that the spectrum of $g_{J,j}$ is contained in $[n_{J,j}, m_{J,j}]$ with

$$\begin{cases} n_{J,j} &= (2J-1)2^j - (2^j - 1) \\ m_{J,j} &= (2J-1)2^j + (2^j - 1). \end{cases}$$

Thus, the spectra of the $g_{J,j}$, $1 \leq J \leq j$ are disjoint. Moreover, $\|g_{J,j}\|_p = 1$ and for $1 \leq J < j$, $\|g_{J,j}\|_p \leq \|\chi_{J,j}\|_p \leq 2^{(J-j+1)/p}$.

We finally set

$$g_j = \sum_{J=1}^j c_{J,j}g_{J,j}$$

and we claim that g_j is the trigonometric polynomial we are looking for. First of all, the spectrum of g_j is included in $[n_{1,j}, m_{j,j}]$ which is contained in $[0, j2^{j+1})$. Moreover, the L^p norm of g_j is

$$\|g_j\|_p \leq \sum_{J=1}^j \frac{1}{j}2^{-(J-j+1)/p}\|g_{J,j}\|_p \leq 1.$$

Pick now any $x \in \mathbf{I}_{J,j}$, $1 \leq J \leq j$ so that

$$\begin{aligned} |S_{m_{J,j}}g_j(x) - S_{n_{J,j}-1}g_j(x)| &= |c_{J,j}g_{J,j}(x)| \\ &= \frac{1}{j}2^{-(J-j+1)/p}|\sigma_{2^j}\chi_{J,j}(x)|. \end{aligned}$$

Observing that $\chi_{J,j}(x) = 1$ and applying the first point of Lemma 1.7 to $1 - \chi_{J,j}$, we find

$$|\sigma_{2^j}\chi_{J,j}(x)| \geq 1 - |\sigma_{2^j}(1 - \chi_{J,j}(x))| \geq \frac{1}{4}.$$

Thus,

$$|S_{m_{J,j}}g_j(x) - S_{n_{J,j}-1}g_j(x)| \geq \frac{1}{4j}2^{-(J-j+1)/p}$$

and the conclusion follows with $n_2 = m_{J,j}$ and $n_1 = n_{J,j} - 1$. □

We are now ready to construct the saturating function. It is defined by

$$g = \sum_{j \geq 1} \frac{1}{j^2} e_{j2^{j+1}} g_j.$$

Observe in particular that the functions $e_{j2^{j+1}} g_j$ have disjoint spectra (the spectrum of $e_{j2^{j+1}} g_j$ is contained in $[j2^{j+1}; j2^{j+2})$) and that g belongs to $L^p(\mathbb{T})$.

We then show that for any $x \in D_\alpha$, $\alpha > 1$,

$$\limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n} \geq \frac{1}{p} \left(1 - \frac{1}{\alpha} \right).$$

Indeed, let $x \in D_\alpha$ and let $\varepsilon > 0$ with $\alpha - \varepsilon > 1$. We can find integers K and J with J as large as we want and $K \notin 2\mathbb{Z}$ such that

$$\left| x - \frac{K}{2^J} \right| \leq \frac{1}{2^{(\alpha-\varepsilon/2)J}}.$$

We set $j = [(\alpha - \varepsilon/2)J]$ the integer part of $(\alpha - \varepsilon/2)J$ and k such that $k/2^j = K/2^J$. Hence,

$$\left| x - \frac{k}{2^j} \right| \leq \frac{1}{2^{(\alpha-\varepsilon/2)J}} \leq \frac{1}{2^j}.$$

Using Lemma 2.2, we can find two integers n_1 and n_2 satisfying $j2^{j+1} \leq n_1 < n_2 < j2^{j+2}$ and such that

$$\begin{aligned} |S_{n_2} g(x) - S_{n_1} g(x)| &= \frac{1}{j^2} |S_{n_2}(e_{j2^{j+1}} g_j)(x) - S_{n_1}(e_{j2^{j+1}} g_j)(x)| \\ &\geq \frac{1}{4j^3} 2^{-(J-j+1)/p} \\ &\geq \frac{1}{4j^3} 2^{\frac{1}{p}(j - \frac{j+1}{\alpha-\varepsilon/2} - 1)} \\ &\geq C 2^{\frac{1}{p}(1 - \frac{1}{\alpha-\varepsilon})j}. \end{aligned}$$

It follows that we can find $n \in \{n_1, n_2\}$ such that $|S_n g(x)| \geq \frac{C}{2} 2^{\frac{1}{p}(1 - \frac{1}{\alpha-\varepsilon})j}$. Combining the estimates on n and on $|S_n g(x)|$, and since J (hence j , hence n) can be taken as large as we want, we get that

$$\limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n} \geq \frac{1}{p} \left(1 - \frac{1}{\alpha - \varepsilon} \right).$$

Since $\varepsilon > 0$ is arbitrary, we obtain in fact that

$$\text{for any } x \in D_\alpha, \quad \limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n} \geq \frac{1}{p} \left(1 - \frac{1}{\alpha} \right).$$

At this point, it would be nice to get a lower bound for $\limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n}$ for any x with dyadic exponent equal to α . Unfortunately, this does not seem easy and we will rather

conclude by using an argument lying on Hausdorff measures. Indeed, define

$$\begin{aligned} D_\alpha^1 &= \left\{ x \in D^\alpha; \limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n} = \frac{1}{p} \left(1 - \frac{1}{\alpha} \right) \right\} \\ D_\alpha^2 &= \left\{ x \in D^\alpha; \limsup_{n \rightarrow +\infty} \frac{\log |S_n g(x)|}{\log n} > \frac{1}{p} \left(1 - \frac{1}{\alpha} \right) \right\}. \end{aligned}$$

We have already observed that $\mathcal{H}^{1/\alpha}(D_\alpha^1 \cup D_\alpha^2) = \mathcal{H}^{1/\alpha}(D_\alpha) = +\infty$. It suffices to prove that $\mathcal{H}^{1/\alpha}(D_\alpha^2) = 0$. Let (β_n) be a sequence of real numbers such that $\beta_n > \frac{1}{p} \left(1 - \frac{1}{\alpha} \right)$

and $\lim_{n \rightarrow +\infty} \beta_n = \frac{1}{p} \left(1 - \frac{1}{\alpha} \right)$.

Let us observe that

$$D_\alpha^2 \subset \bigcup_{n \geq 0} \mathcal{E}(\beta_n, g).$$

Moreover, Theorem 1.1 implies that $\mathcal{H}^{1/\alpha}(\mathcal{E}(\beta_n, g)) = 0$ for all n . Hence, $\mathcal{H}^{1/\alpha}(D_\alpha^2) = 0$ and $\mathcal{H}^{1/\alpha}(D_\alpha^1) = +\infty$, which proves that

$$\dim_{\mathcal{H}} \left(E \left(\frac{1}{p} \left(1 - \frac{1}{\alpha} \right), g \right) \right) \geq \frac{1}{\alpha}.$$

By Theorem 1.1 again, this inequality is necessarily an equality. Finally, g satisfies the conclusions of Theorem 1.2, setting $1 - \beta p = 1/\alpha$.

Remark 2.3. If $\alpha = 1$, then $\beta = 0$ and the conclusion is a consequence of Carleson's Theorem.

2.2. The residual set. To build the dense G_δ -set, the idea is that any function whose Fourier coefficients are sufficiently close to those of the saturating function g on infinitely many intervals $[j2^{j+1}, j2^{j+2})$ will satisfy the conclusions of Theorem 1.2. Precisely, let $(f_j)_{j \geq 1}$ be a dense sequence of polynomials in $L^p(\mathbb{T})$ with spectrum contained in $[-j, j]$. We define a sequence $(h_j)_{j \geq 1}$ as follows:

$$h_j = f_j + \frac{1}{j} e_{j2^{j+1}} g_j$$

so that $\|h_j - f_j\|_p$ goes to 0 and $(h_j)_{j \geq 1}$ remains dense in $L^p(\mathbb{T})$. Observe also that the spectra of f_j and $h_j - f_j$ don't overlap. Finally, let $(r_j)_{j \geq 1}$ be a sequence of positive integers so small that, for any $f \in L^p(\mathbb{T})$ with $\|f\|_{L^p} \leq r_j$, $\|S_n f\|_\infty \leq 1$ for any $n \leq j2^{j+2}$. The dense G_δ set we will consider is

$$A = \bigcap_{l \in \mathbb{N}} \bigcup_{j \geq l} B(h_j, r_j).$$

Let f belong to A and let $(j_l)_{l \geq 1}$ be an increasing sequence of integers such that f belongs to $B(h_{j_l}, r_{j_l})$ for any l . Then, for any $\alpha > 1$, we define $J_l = [j_l/\alpha] + 1$ (which is smaller than j_l if l is large enough) and

$$E = \limsup_{l \rightarrow +\infty} \mathbf{I}_{J_l, j_l}.$$

For any $x \in E$ one can find $j = j_l$ as large as we want, the corresponding $J = J_l$ and $1 \leq k \leq 2^j - 1$ such that x belongs to $I_{k,j}$ with $k/2^j \in \mathcal{I}_J$.

Observe that $f = f_j + \frac{1}{j}e_{j2^{j+1}}g_j + (f - h_j)$. By Lemma 2.2, we can find two integer n_1 and n_2 satisfying $j2^{j+1} \leq n_1 < n_2 < j2^{j+2}$ and such that

$$|S_{n_2}(e_{j2^{j+1}}g_j)(x) - S_{n_1}(e_{j2^{j+1}}g_j)(x)| \geq \frac{1}{4j}2^{-(J-j+1)/p}.$$

Using the definition of the r_j , we obtain

$$\begin{aligned} |S_{n_2}f(x) - S_{n_1}f(x)| &\geq \frac{1}{4j^2}2^{-(J-j+1)/p} - |S_{n_2}(f - h_j)(x)| - |S_{n_1}(f - h_j)(x)| \\ &\geq \frac{1}{4j^2}2^{-(J-j+1)/p} - 2 \end{aligned}$$

so that

$$|S_{n_2}f(x)| \geq \frac{C}{j^2}2^{-(J-j+1)/p} \quad \text{or} \quad |S_{n_1}f(x)| \geq \frac{C}{j^2}2^{-(J-j+1)/p}.$$

Observing that

$$\begin{cases} \max(\log n_2, \log n_1) &= j \log 2 + O(\log j) \\ \log(j^{-2}2^{-(J-j+1)/p}) &= \frac{1}{p}(1 - \frac{1}{\alpha})j \log 2 + O(\log j) \end{cases}$$

we find in particular that, for any $x \in E$,

$$\limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log n} \geq \frac{1}{p} \left(1 - \frac{1}{\alpha}\right).$$

On the other hand, let us write

$$\mathbf{I}_{J_l, j_l} = \bigcup_{1 \leq K < 2^{J_l}, K \notin 2\mathbb{Z}} \left[\frac{K}{2^{J_l}} - \frac{1}{2^{j_l}}, \frac{K}{2^{J_l}} + \frac{1}{2^{j_l}} \right]$$

and remark that for any l , since $J_l \geq j_l/\alpha$,

$$\bigcup_{1 \leq K < 2^{J_l}, K \notin 2\mathbb{Z}} \left[\frac{K}{2^{J_l}} - \frac{1}{2^{j_l/\alpha}}, \frac{K}{2^{J_l}} + \frac{1}{2^{j_l/\alpha}} \right] \supset [0, 1].$$

Hence, we can apply Corollary 1.9 to get $\mathcal{H}^{1/\alpha}(E) = +\infty$. We now conclude exactly as in Section 2.1 to get $\mathcal{H}^{1/\alpha}(E^1) = +\infty$, with

$$E^1 = \left\{ x \in E; \limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log n} = \frac{1}{p} \left(1 - \frac{1}{\alpha}\right) \right\}.$$

Finally,

$$\dim_{\mathcal{H}} \left(E \left(\frac{1}{p} \left(1 - \frac{1}{\alpha}\right), f \right) \right) \geq \frac{1}{\alpha}$$

and f satisfies the conclusions of Theorem 1.2, setting $1 - \beta p = 1/\alpha$.

Remark 2.4. During the construction, we didn't use that the spectra of the functions $e_{j2^{j+1}}g_j$ are disjoint, because we considered each one separately. We could also define h_j by $h_j = f_j + \frac{1}{j}e_{j+1}g_j$.

Remark 2.5. The above construction can be carried on $L^1(\mathbb{T})$. Namely, for quasi-all $f \in L^1(\mathbb{T})$, we obtain for any $\beta \in [0, 1]$,

$$\dim_{\mathcal{H}}(E(\beta, f)) \geq 1 - \beta.$$

However, we cannot go further because Carleson's Theorem dramatically breaks down in $L^1(\mathbb{T})$ and we do not have Theorem 1.1 at our disposal in this context. The study of what happens exactly on $L^1(\mathbb{T})$ is a very exciting open question.

3. MULTIFRACTAL ANALYSIS OF THE DIVERGENCE OF THE FOURIER SERIES OF FUNCTIONS OF $\mathcal{C}(\mathbb{T})$

We turn to the proof of Theorem 1.6. We follow a strategy close to that of Section 2. First of all, we will give an upper bound for the precised Hausdorff dimension of the sets $\mathcal{F}(\beta, f)$ (hence, of the sets $F(\beta, f)$) for any $f \in \mathcal{C}(\mathbb{T})$ and any $\beta \in (0, 1)$. Second, we will build polynomials with small L^∞ -norms and such that their Fourier series have big partial sums on big intervals. These polynomials will be the blocks of our final construction. Working on $\mathcal{C}(\mathbb{T})$ adds several difficulties which will be explained when we will encounter them.

3.1. The sets $\mathcal{F}(\beta, f)$ cannot be too big. We shall prove the following lemma (recall that $\phi_{s,t}(x) = x^s \exp((\log 1/x)^{1-t})$).

Lemma 3.1. *Let $\beta \in (0, 1)$ and $f \in \mathcal{C}(\mathbb{T})$. Then, for any $\gamma > 1 - \beta$,*

$$\mathcal{H}^{\phi_{1,\gamma}}(\mathcal{F}(\beta, f)) = 0.$$

In particular, the precised Hausdorff dimension of $\mathcal{F}(\beta, f)$ cannot exceed $(1, 1 - \beta)$.

Proof. A key point in Aubry's proof of Theorem 1.1 is the Carleson-Hunt theorem which asserts that, for any $g \in L^p(\mathbb{T})$, $1 < p < +\infty$,

$$\|S^*g\|_p \leq C_p \|g\|_p \quad \text{where} \quad S^*g(x) = \sup_{n \geq 0} |S_n g(x)|.$$

On $\mathcal{C}(\mathbb{T})$, a weak inequality (also due to Hunt) remains valid (see [1, Theorem 12.5]): there are two absolute constants $A, B > 0$ such that, for every $f \in \mathcal{C}(\mathbb{T})$ and every $y > 0$,

$$\lambda(\{x \in \mathbb{T}; S^*f(x) > y\}) \leq Ae^{-By/\|f\|_\infty}.$$

Here, λ denotes the Lebesgue measure on \mathbb{T} .

So, let $\beta \in (0, 1)$ and $f \in \mathcal{C}(\mathbb{T})$. We may assume $\|f\|_\infty \leq 1$. For any $M > 0$, we introduce

$$\mathcal{F}(\beta, f, M) = \left\{ x \in \mathbb{T}; \limsup_{n \rightarrow +\infty} (\log n)^{-\beta} |S_n f(x)| > M \right\}.$$

Since $\mathcal{F}(\beta, f) = \bigcup_{M > 0} \mathcal{F}(\beta, f, M)$, we just need to prove that $\mathcal{H}^{\phi_{1,\gamma}}(\mathcal{F}(\beta, f, M)) = 0$ for every $M > 0$. From now on, we fix some $M > 0$. We pick any $x \in \mathcal{F}(\beta, f, M)$ and n_x large enough such that

$$|S_{n_x} f(x)| \geq M(\log n_x)^\beta.$$

Such an inequality remains true in an interval around x whose size is not so small. Precisely, because n_x can be assumed to be large and since the L^1 -norm of the Dirichlet

kernel D_n behaves like $\frac{4}{\pi^2} \log n$, we may assume that $\|S_{n_x} f\|_\infty \leq (\log n_x) \|f\|_\infty \leq \log n_x$. By Bernstein's inequality, $\|(S_{n_x} f)'\|_\infty \leq n_x \log n_x$. Let

$$I_x = \left[x - \frac{M}{2n_x(\log n_x)^{1-\beta}}, x + \frac{M}{2n_x(\log n_x)^{1-\beta}} \right].$$

For any $y \in I_x$, we get

$$(3) \quad |S_{n_x} f(y)| \geq \frac{M}{2} (\log n_x)^\beta.$$

$(I_x)_{x \in \mathcal{F}(\beta, f, M)}$ is a covering of $\mathcal{F}(\beta, f, M)$. We can extract a Vitali's covering, namely a countable family of disjoint intervals $I_i, i \in \mathbb{N}$, of length $\frac{M}{n_i(\log n_i)^{1-\beta}}$ such that $\mathcal{F}(\beta, f, M) \subset \bigcup_i 5I_i$. Let us finally set, for any $q \geq 1$, $\mathcal{U}_q = \left\{ i; 2^{q+1} \geq \frac{M(\log n_i)^\beta}{2} > 2^q \right\}$. Without loss of generality, we may assume the n_i so large that $\bigcup_q \mathcal{U}_q = \mathbb{N}$. By applying Hunt's theorem,

$$\lambda(\{x; S^* f(x) > 2^q\}) \leq Ae^{-B2^q}.$$

Now, by (3), the set $\{x; S^* f(x) > 2^q\}$ contains the disjoint intervals I_i , for $i \in \mathcal{U}_q$. Thus,

$$\sum_{i \in \mathcal{U}_q} |I_i| \leq Ae^{-B2^q}.$$

Moreover, for any $i \in \mathcal{U}_q$, it is not hard to check that

$$|I_i| \geq Ce^{-D2^{q/\beta}}$$

for some positive constants C, D which do not depend on q . Picking any α such that $1 - \beta < \alpha < \gamma$, we get

$$\begin{aligned} \sum_{i \in \mathcal{U}_q} \phi_{1,\alpha}(5|I_i|) &= \sum_{i \in \mathcal{U}_q} 5|I_i| \exp((\log(1/5|I_i|))^{1-\alpha}) \\ &\leq 5 \left(\sum_{i \in \mathcal{U}_q} |I_i| \right) \exp\left(\left(D2^{q/\beta} - \log 5C\right)^{1-\alpha}\right) \\ &\leq 5Ae^{-B2^q + D'2^{q(1-\alpha)/\beta}}. \end{aligned}$$

Since $1 - \alpha < \beta$, this shows that there exists $C_0 < +\infty$ such that

$$\sum_{i \in \mathbb{N}} \phi_{1,\alpha}(5|I_i|) = \sum_{q \in \mathbb{N}} \sum_{i \in \mathcal{U}_q} \phi_{1,\alpha}(5|I_i|) \leq C_0.$$

Remember that $\bigcup_i 5I_i$ is a covering of $\mathcal{F}(\beta, f, M)$ and that the I_i can be chosen as small as we want. We can then conclude that $\mathcal{H}^{\phi_{1,\alpha}}(\mathcal{F}(\beta, f, M)) \leq C_0$. In particular, $\mathcal{H}^{\phi_{1,\gamma}}(\mathcal{F}(\beta, f, M)) = 0$, since $\phi_{1,\alpha} \gg \phi_{1,\gamma}$. \square

Remark 3.2. The functions $\phi_{1,\gamma}$, for $\gamma > 1 - \beta$, are not optimal in the statement of the previous lemma. We can replace them by any function $\phi(x) = x (\exp((\log 1/x)^\beta \varepsilon(x)))$ with $\varepsilon(x)$ goes to 0 as x goes to 0.

3.2. The basic construction. When we try to build explicitly a continuous function whose Fourier series diverges at some point, say 0, a natural way is to consider polynomials P with small L^∞ norm, and satisfying nevertheless that $|S_n P(0)|$ is big for some large value of n . The easiest examples are

$$P_N(x) = e_N(x) \sum_{j=1}^N \frac{\sin(2\pi jx)}{j},$$

since the sequence $(\|P_N\|_\infty)_{N \geq 1}$ is bounded whereas $|S_N(P)(0)| \sim \frac{1}{2} \log N$. Moreover, this example is in some sense optimal since $\|S_N f\|_\infty \leq C(\log N)\|f\|_\infty$ for any $f \in \mathcal{C}(\mathbb{T})$.

In our context, we need to find a polynomial P which satisfies a similar property not only at one point, but on a set which is rather big since at the end we want to construct sets of divergence with Hausdorff dimension 1. This does not seem to be the case for P_N , the reason being that $|(S_N P)'(0)|$ behaves like N , which is much bigger than $S_N P(0)$.

To tackle this problem, we start from a construction of Kahane and Katznelson in [8] (see also [9]) which they use to prove that every subset of \mathbb{T} of Lebesgue measure 0 is a set of divergence for $\mathcal{C}(\mathbb{T})$. Since we want to control both the size of the sets E and the index n such that $S_n P(x)$ becomes larger than some given real number for any $x \in E$, the forthcoming lemma needs very careful estimations.

Lemma 3.3. *Let $\beta \in (0, 1)$, $\delta \in (0, 1)$ and $K \geq 2$. Then there exist an integer $k \geq K$, an integer n as large as we want and a trigonometric polynomial P with spectrum contained in $[0, 2n - 1]$ such that*

- $|P(x)| \leq 1$ for any $x \in \mathbb{T}$;
- $\log |S_n P(x)| \geq (1 - \delta)\beta \log \log n$ for any $x \in I_k^\beta$,

where $I_k^\beta = \bigcup_{j=0}^{k-1} \left[\frac{j}{k} - \frac{1}{2k \exp((\log k)^\beta)}; \frac{j}{k} + \frac{1}{2k \exp((\log k)^\beta)} \right]$.

Proof. Let us first describe the idea of the proof. We shall construct a trigonometric polynomial Q with spectrum in $[1, n - 1]$ and with the following properties: $|\Im Q|$ is small and $|Q|$ is large on a set E . We then set $P = e_n \times \Im Q$, so that $\|P\|_\infty$ is small. On the other hand, writing $Q = \sum_{k=1}^{n-1} a_k e_k$, $2i \Im Q = -\sum_{k=1}^{n-1} \bar{a}_k e_{-k} + \sum_{k=1}^{n-1} a_k e_k$, so that

$$|S_n(P)| = \frac{1}{2} \left| \sum_{k=1}^{n-1} \bar{a}_k e_{n-k} \right| = \frac{1}{2} \left| \sum_{k=1}^{n-1} a_k e_k \right| = \frac{1}{2} |Q|$$

is large on E . The construction of Q will be done by taking $\log f$, the logarithm of an holomorphic function defined on a neighbourhood of the closed unit disk $\overline{\mathbb{D}}$ (which allows to control the imaginary part of $\log f$ while the modulus of it can be large), and by taking a Fejér sum of $\log f$.

We now proceed with the details. The proof uses holomorphic functions and it is better to see \mathbb{T} as the boundary of the unit disk \mathbb{D} . To avoid cumbersome notations, the letter C will denote throughout the proof a positive and absolute constant, whose value may

change from line to line. Let $k \geq K$ whose value will be fixed later. We set:

$$\begin{aligned}\varepsilon &= \frac{1}{k \exp((\log k)^\beta)} \\ z_j &= e^{\frac{2\pi ij}{k}}, \quad j = 0, \dots, k-1 \\ f(z) &= \frac{1}{k} \sum_{j=0}^{k-1} \frac{1+\varepsilon}{1+\varepsilon - \bar{z}_j z}.\end{aligned}$$

f is holomorphic in a neighbourhood of $\bar{\mathbb{D}}$. We claim that f satisfies the following four properties.

- (P1): $\forall z \in \bar{\mathbb{D}}, \quad \Re f(z) \geq C\varepsilon;$
- (P2): $\forall z \in I_k^\beta, \quad |f(z)| \geq \Re f(z) \geq C \exp((\log k)^\beta);$
- (P3): $\forall z \in \mathbb{T}, \quad |f(z)| \leq C \exp((\log k)^\beta);$
- (P4): $\forall z \in \mathbb{T}, \quad \left| \frac{f'(z)}{f(z)} \right| \leq \frac{C}{\varepsilon^3}.$

Indeed, for any $z \in \bar{\mathbb{D}}$ and any $j \in \{0, \dots, k-1\}$,

$$(4) \quad \Re \left(\frac{1+\varepsilon}{1+\varepsilon - \bar{z}_j z} \right) = \frac{1+\varepsilon}{|1+\varepsilon - \bar{z}_j z|^2} \Re(1+\varepsilon - z_j \bar{z}) \geq \frac{1+\varepsilon}{(2+\varepsilon)^2} \times \varepsilon \geq C\varepsilon,$$

which proves (P1). To prove (P2), we may assume that $z = e^{2\pi i\theta}$ with $\theta \in [\frac{-\varepsilon}{2}; \frac{\varepsilon}{2}]$. Then

$$\Re \left(\frac{1+\varepsilon}{1+\varepsilon - \bar{z}_0 z} \right) = \frac{1+\varepsilon}{|1+\varepsilon - z|^2} \Re(1+\varepsilon - \bar{z}) \geq \frac{C}{\varepsilon}.$$

If we combine this with (4), we get

$$\Re f(z) \geq \frac{C}{k\varepsilon} + \frac{k-1}{k} C\varepsilon \geq \frac{C}{k\varepsilon} = C \exp((\log k)^\beta).$$

which gives (P2).

Conversely, we want to control $\sup_{z \in \mathbb{T}} |f(z)|$. Pick any $z = e^{2\pi i\theta} \in \mathbb{T}$. By symmetry, we may and shall assume that $|\theta| \leq \frac{1}{2k}$. Then we get

$$\left| \frac{1+\varepsilon}{1+\varepsilon - \bar{z}_0 z} \right| \leq \frac{C}{\varepsilon}.$$

Now, for any $j \in \{1, \dots, k/4\}$, we can write

$$\begin{aligned}|1+\varepsilon - \bar{z}_j z| &\geq |\Im m(\bar{z}_j z)| \\ &\geq \sin \left(\frac{2\pi j}{k} - 2\pi\theta \right) \\ &\geq \frac{2}{\pi} \times 2\pi \left(\frac{j}{k} - \theta \right) \\ &\geq \frac{4}{k} \left(j - \frac{1}{2} \right).\end{aligned}$$

Taking the sum,

$$\left| \sum_{j=1}^{k/4} \frac{1+\varepsilon}{1+\varepsilon - \bar{z}_j z} \right| \leq \frac{k(1+\varepsilon)}{4} \sum_{j=1}^{k/4} \frac{1}{j-1/2} \leq Ck \log k.$$

In the same way, we obtain

$$\left| \sum_{j=3k/4}^{k-1} \frac{1+\varepsilon}{1+\varepsilon-\bar{z}_j z} \right| \leq Ck \log k$$

whereas $|1+\varepsilon-\bar{z}_j z| \geq C$ for any $j \in [k/4, 3k/4]$, so that

$$\left| \sum_{j=k/4}^{3k/4} \frac{1+\varepsilon}{1+\varepsilon-\bar{z}_j z} \right| \leq Ck.$$

Putting this together, we get

$$|f(z)| = \left| \frac{1}{k} \sum_{j=0}^{k-1} \frac{1+\varepsilon}{1+\varepsilon-\bar{z}_j z} \right| \leq C \left(\frac{1}{k\varepsilon} + \log k + 1 \right) \leq C \exp((\log k)^\beta).$$

Finally, it remains to prove **(P4)**. We observe that

$$f'(z) = \frac{1}{k} \sum_{j=0}^{k-1} \frac{(1+\varepsilon)\bar{z}_j}{(1+\varepsilon-\bar{z}_j z)^2}.$$

We do not try to get a very precise estimate for $|f'(z)|$ (this is not useful for us). We just observe that $|1+\varepsilon-\bar{z}_j z|^2 \geq \varepsilon^2$ for any $j \in \{0, \dots, k-1\}$ and any $z \in \mathbb{T}$, so that

$$|f'(z)| \leq \frac{C}{\varepsilon^2}.$$

If we combine this with **(P1)**, we get **(P4)**.

We are almost ready to construct P . The next step is to take $h(z) = \log(f(z))$, which defines a holomorphic function in a neighbourhood of $\bar{\mathbb{D}}$ by **(P1)**. Moreover, $|\Im m(h(z))| \leq \pi/2$ for any $z \in \bar{\mathbb{D}}$ and $h(0) = 0$. Now, we look at the function h on the boundary of the unit disk \mathbb{D} , that is we introduce the function $g(x) = h(e^{2i\pi x})$ defined on the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. Properties **(P2)**, **(P3)** and **(P4)** can be rewritten as

$$\begin{aligned} \forall x \in I_k^\beta, \quad |g(x)| &\geq (\log k)^\beta - C \\ \forall x \in \mathbb{T}, \quad |g(x)| &\leq (\log k)^\beta + C \\ \forall x \in \mathbb{T}, \quad |g'(x)| &\leq Ck^3 \exp(3(\log k)^\beta). \end{aligned}$$

Let now n be the smallest integer such that $Ck^3 \exp(3(\log k)^\beta) \leq n$. We also have $\|g'\|_\infty \leq n$ and we can apply the second part of Lemma 1.7 to the function $\theta(t) = g(t) - g(x)$ when $x \in I_k^\beta$. Recall that $\|\theta\|_\infty \leq 2(\log k)^\beta + C$. We get

$$|\sigma_n \theta(x)| \leq \frac{(\log k)^\beta}{2} + C$$

and we can conclude that

$$|\sigma_n g(x)| \geq |g(x)| - |\sigma_n \theta(x)| \geq \frac{(\log k)^\beta}{2} - C.$$

We finally set

$$P = \frac{2}{\pi} e_n \sigma_n (\Im m g) = \frac{2}{\pi} e_n \Im m (\sigma_n g).$$

It is straightforward to check that $\|P\|_\infty \leq 1$ (recall that σ_n is a contraction on $\mathcal{C}(\mathbb{T})$), and that the spectrum of $\sigma_n g$ is contained in $[1, n-1]$ ($\hat{g}(0) = 0$ since $h(0) = 0$). Now, the simple algebraic trick exposed at the beginning of the proof shows that

$$|S_n P(x)| = \left| \frac{1}{\pi} \sigma_n g(x) \right|,$$

so that, for any $x \in I_k^\beta$,

$$|S_n P(x)| \geq \frac{1}{2\pi} (\log k)^\beta - C.$$

This leads to

$$\log |S_n P(x)| \geq \beta \log \log k - C.$$

On the other hand,

$$\begin{aligned} \log \log n &\leq \log \left(3 \log k + 3(\log k)^\beta + \log C \right) \\ &\leq \log \log k + C. \end{aligned}$$

Finally,

$$\frac{\log \log |S_n P(x)|}{\log \log n} \geq \frac{\beta \log \log k - C}{\log \log k + C} \geq (1 - \delta)\beta,$$

provided k has been chosen large enough. Moreover, n can be chosen as large as we want since $n \rightarrow +\infty$ when $k \rightarrow +\infty$. \square

Remark 3.4. The fact that we have to compare $\log \log n$ and $\log |S_n|$ helps us for the previous proof. Even if n and k do not have the same order of growth, this is not apparent when we apply the iterated logarithm.

Remark 3.5. During the construction, the integers k and n can't be chosen independently : they satisfy $n - 1 \leq Ck^3 \exp(3(\log k)^\beta) \leq n$ where C is an absolute constant. If we want to construct a polynomial P satisfying the conclusion of Lemma 3.3 with a large value of n , we need also to choose a large value of k .

3.3. The conclusion. We are now going to prove the full statement of Theorem 1.6. At this point, the situation is less favourable than in the L^p -case. There, the basic construction done at each step j did not depend on the index of divergence that we would like to get. We had the same function g_j which worked for all indices of divergence, and it was the dyadic exponent of x which decided how large was $|g_j(x)|$. The construction done in Lemma 3.3 is less efficient, because the polynomial P does depend on the expected divergence index β (the index β is a parameter of the definition of f above). We have to overcome this new difficulty and the solution will be to introduce redundancy in the construction of the G_δ -set.

As usual, we start from a sequence $(f_j)_{j \geq 1}$ of polynomials which is dense in $\mathcal{C}(\mathbb{T})$. For convenience, we assume that $\|f_j\|_\infty \leq j$ for any j and that the spectrum of f_j is contained in $[-j, j]$. Furthermore, we fix four sequences (α_l) , (β_l) , (δ_l) and (ε_l) with values in $(0, 1)$ and such that:

- (β_l) is dense in $(0, 1)$ and $l \mapsto \beta_l$ is one to one;
- $\sum_l \varepsilon_l \leq 1$;
- (δ_l) and (α_l) go to zero.

- $\delta_l < 1/3$.

Let now $j \geq 1$. By induction on $l = 1, \dots, j$, we build sequences $(P_{j,l})$, $(n_{j,l})$ and $(k_{j,l})$ satisfying the conclusions of Lemma 3.3 with $\beta = \beta_l$, $\delta = \delta_l$ and $K = j$ (to ensure that $\lim_{j \rightarrow +\infty} k_{j,l} = +\infty$) and we will decide how large should be $n_{j,l}$ during the construction. According to Remark 3.5, these constraints on $n_{j,l}$ will determine the values of the $k_{j,l}$. We then set

$$g_j := f_j + \alpha_j \sum_{l=1}^j \varepsilon_l e_{n_{j,l}} P_{j,l}$$

so that $\|g_j - f_j\|_\infty \leq \alpha_j \sum_{l=1}^j \varepsilon_l \|P_{j,l}\|_\infty \leq \alpha_j$. In particular, the sequence (g_j) remains dense in $\mathcal{C}(\mathbb{T})$.

Recall that the spectrum of f_j is included in $[-j, j]$ and observe that the spectrum of $e_{n_{j,l}} P_{j,l}$ lies in $[n_{j,l}, 3n_{j,l} - 1]$. If we suppose that $n_{j,1} = j + 1$ and $n_{j,l+1} \geq 3n_{j,l}$, we can conclude that the spectra of $f_j, e_{n_{j,1}} P_{j,1}, \dots, e_{n_{j,j}} P_{j,j}$ are disjoint.

Let now x belongs to $I_{k_{j,l}}^{\beta_l}$ for some $l \leq j$. Then

$$\begin{aligned} |S_{2n_{j,l}} g_j(x)| &\geq \alpha_j \varepsilon_l |S_{n_{j,l}} P_{j,l}(x)| - \alpha_j \sum_{m=1}^{l-1} \varepsilon_m \|P_{j,m}\|_\infty - j \\ &\geq \alpha_j \varepsilon_l |S_{n_{j,l}} P_{j,l}(x)| - \alpha_j - j. \end{aligned}$$

Because we can choose $n_{j,l}$ as large as we want in the process, we may always assume that the choice that we have done ensures that

$$|S_{2n_{j,l}} g_j(x)| \geq \frac{\alpha_j \varepsilon_l}{2} |S_{n_{j,l}} P_{j,l}(x)|.$$

Taking the logarithm, we find

$$\begin{aligned} \log |S_{2n_{j,l}} g_j(x)| &\geq \log |S_{n_{j,l}} P_{j,l}(x)| + \log \varepsilon_l + \log \alpha_j - \log 2 \\ &\geq (1 - \delta_l) \beta_l \log \log(n_{j,l}) + \log \varepsilon_l + \log \alpha_j - \log 2 \\ &\geq (1 - 2\delta_l) \beta_l \log \log(2n_{j,l}) \end{aligned}$$

provided again that we have chosen $n_{j,l}$ very large.

We then fix $r_j > 0$ so small that, for any $f \in B(g_j, r_j)$ (the balls are related to the norm $\|\cdot\|_\infty$), for any $l \leq j$,

$$\|S_{2n_{j,l}} f - S_{2n_{j,l}} g_j\|_\infty \leq 1/2.$$

Observe that for every real number $t \geq 1$, we have $\log(t - 1/2) \geq \log(t) - \log 2$. For any $x \in I_{k_{j,l}}^{\beta_l}$ with $l \leq j$, we get

$$\begin{aligned} \log |S_{2n_{j,l}} f(x)| &\geq \log |S_{2n_{j,l}} g_j(x)| - \log 2 \\ &\geq (1 - 2\delta_l) \beta_l \log \log(2n_{j,l}) - \log 2 \\ &\geq (1 - 3\delta_l) \beta_l \log \log(2n_{j,l}) \end{aligned}$$

if $n_{j,l}$ are chosen sufficiently large such that $\delta_l \beta_l \log \log(2n_{j,l}) \geq \log 2$.

We finally set

$$A = \bigcap_{p \in \mathbb{N}} \bigcup_{j \geq p} B(g_j, r_j),$$

and we claim that A is the dense G_δ set we are looking for.

Indeed, let f belong to A and let (j_p) be an increasing sequence of integers such that for every $p \geq 0$, $f \in B(g_{j_p}, r_{j_p})$. We consider $\beta \in (0, 1)$ and choose p_0 such that

$$\{\beta_1, \dots, \beta_{j_{p_0}}\} \cap (0, \beta) \neq \emptyset.$$

Such a p_0 exists since the sequence $(\beta_l)_{l \geq 1}$ is dense in $(0, 1)$. For every $p \geq p_0$, let l_p be chosen in $\{1, \dots, j_p\}$ such that

$$\beta - \beta_{l_p} = \inf\{\beta - \beta_l; l \leq j_p \text{ and } \beta > \beta_l\}.$$

Since the sequence (β_l) is dense in $(0, 1)$, $\beta_{l_p} < \beta$ for $p \geq p_0$ and $\beta_{l_p} \rightarrow \beta$. Moreover, since $l \mapsto \beta_l$ is one to one, it is clear that l_p is non decreasing and goes to $+\infty$.

Observe that, for $p \geq p_0$, $I_{k_{j_p}, l_p}^\beta \subset I_{k_{j_p}, l_p}^{\beta_{l_p}}$, so that, for any $x \in I_{k_{j_p}, l_p}^\beta$, setting $N_p = 2n_{j_p, l_p}$,

$$\log |S_{N_p} f(x)| \geq (1 - 3\delta_{l_p})\beta_{l_p} \log \log(N_p).$$

In particular, setting $F = \limsup_p I_{k_{j_p}, l_p}^\beta$, we get that

$$\limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log \log n} \geq \beta$$

for any $x \in F$. Now, we can apply Corollary 1.9 with a gauge function ϕ satisfying $\phi^{-1}(y) = y \exp[-(\log(1/2y))^\beta]$ to obtain $\mathcal{H}^\phi(F) = \infty$.

Observe that if $y = \phi(x)$, then

$$y = x \exp\left[(\log(1/2y))^\beta\right] \quad \text{and} \quad \log x \leq \log y.$$

It follows that $\phi(x) \leq x \exp[(\log(1/2x))^\beta] \leq \phi_{1, 1-\beta}(x)$ and $\mathcal{H}^{\phi_{1, 1-\beta}}(F) = +\infty$.

We now conclude exactly as in the L^p -case, using Lemma 3.1 to replace Aubry's result. Namely, we set

$$\begin{aligned} F^1 &= \left\{ x \in F; \limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log \log n} = \beta \right\} \\ F^2 &= \left\{ x \in F; \limsup_{n \rightarrow +\infty} \frac{\log |S_n f(x)|}{\log \log n} > \beta \right\} \end{aligned}$$

and we observe that Lemma 3.1 guarantees that $\mathcal{H}^{\phi_{1, 1-\beta}}(F^2) = 0$. Thus, $\mathcal{H}^{\phi_{1, 1-\beta}}(F^1) = +\infty$ and the precised Hausdorff dimension of $F(\beta, f)$, which contains F^1 , is at least $(1, 1 - \beta)$. By Lemma 3.1, it is exactly $(1, 1 - \beta)$.

Remark 3.6. It is amazing that, with our method, it is easier to prove Theorem 1.6 and to deduce Theorem 1.4 from it than to prove Theorem 1.4 directly. Indeed, to ensure that the sets $\mathcal{F}(\beta, f)$ are big, we need to know that the sets $F(\beta', f)$ are small for $\beta' > \beta$. This cannot be done if we stay within the notion of Hausdorff dimension.

Remark 3.7. The method developed above allows us to construct a ‘‘concrete function’’ that satisfies the conclusion of Theorem 1.6. More precisely, it suffices to consider

$$g = \sum_{j=1}^{+\infty} \frac{1}{j^2} \sum_{l=1}^j \varepsilon_l e_{n_{j,l}} P_{j,l}$$

with the constraint $3n_{j,j} < n_{j+1,1}$ to ensure that the blocks $\sum_{l=1}^j \varepsilon_l e_{n_{j,l}} P_{j,l}$ have disjoint spectra. Such a function is some kind of saturating function in the continuous case.

REFERENCES

- [1] J. ARIAS DE REYNA, Pointwise convergence of Fourier series, *Springer Lecture Notes in Math.*, vol 1785.
- [2] J-M. AUBRY, On the rate of pointwise divergence of Fourier and wavelet series in L^p , *Journal of Approx. Th.* **538** (2006), 97-111.
- [3] V. BERESNEVICH, S. VELANI, A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures, *Annals of Math* **164** (2006), 971-992.
- [4] M.M. DODSON, M.V. MELIÁN, D. PESTANA AND S.L. VÉLANI, Patterson measure and Ubiquity, *Ann. Acad. Sci. Fenn.* **20** (1995), 37-60.
- [5] K. FALCONER, Fractal geometry: Mathematical foundations and applications Wiley (2003).
- [6] S. JAFFARD, On the Frisch-Parisi conjecture, *J. Math. Pures Appl.* **79** (2000), 525-552.
- [7] S. JAFFARD, On lacunary wavelet series, *Ann. Appl. Probab.* **10** (2000), 313-329.
- [8] J-P. KAHANE, Y. KATZNELSON, Sur les ensembles de divergence des séries trigonométriques, *Studia Math.* **26** (1966), 305-306.
- [9] Y. KATZNELSON, An Introduction to Harmonic Analysis, John Wiley & Sons, 1968.
- [10] P. MATTILA, Geometry of Sets and Measures in Euclidian Spaces, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1995.
- [11] A. ZYGMUND, Trigonometric series, Cambridge Mathematical Library.

CLERMONT UNIVERSITÉ, UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, BP 10448, F-63000 CLERMONT-FERRAND - CNRS, UMR 6620, LABORATOIRE DE MATHÉMATIQUES, F-63177 AUBIERE

E-mail address: Frederic.Bayart@math.univ-bpclermont.fr, Yanick.Heurteaux@math.univ-bpclermont.fr